
Pr nancy

Pregnancy begins with	and an	Mil ions of micro	oscopic sperm cells		
are formed in the	_ and mix with fluids t	o make	Every month, one egg		
is released from the	and travels throug	gh the fallopian tube	s. If semen gets into		
the during sex, t	he sperm swim up the	cervix into the	and then into		
the fa lopian tubes.					
Sperm can survive	days, and it only tak	ces cel to	get someone		
pregnant. If the sperm reach	nes and fertilizes an eg	g, the egg then divid	les into more cells as		
it moves towards the uterus. If these cells attach to the wall of the uterus, the person					
becomes					
Not interested in getting pre	egnant? It's simple! Op	tion 1 – Don't have	This is the		
only 100% certain way to av	oid pregnancy. Option	2 – f you are having	g sex, use		
control – Condoms, the pill,	an IUD – to reduce yo	ur chances.			
Text: Stay Teen. "Birth Contro 101: Pregnand https://www.youtube.com/watch?v=yqu7Zs/Imaees: "Know Your Body." Wanna Know. A	Ot4Ek&feature=youtu.be. (Accesse	ed October 2, 2019)			




Pr nancy

http://www.iwannaknow.org/teens/sexualhealth/know.html. (Accessed October 1, 2019)

Pregnancy begins with ______ and an ______. Mil ions of microscopic sperm cells are formed in the ______ and mix with fluids to make ______. Every month, one egg is released from the ______ and travels through the fallopian tubes. If semen gets into the ______ during sex, the sperm swim up the cervix into the _____ and then into the fa lopian tubes.

Sperm can survive _____ days, and it only takes _____ cel to get someone pregnant. If the sperm reaches and fertilizes an egg, the egg then divides into more cells as it moves towards the uterus. If these cells attach to the wall of the uterus, the person becomes _____.

Not interested in getting pregnant? It's simple! Option 1 — Don't have ______. This is the only 100% certain way to avoid pregnancy. Option 2 — f you are having sex, use ______ control — Condoms, the pill, an IUD — to reduce your chances.




Sexually Transm tted Diseases (STDs)

contact. Th	at includes genital-to-	contact, which is contact with per	ises, scrotums, vulvas or vaginas, to an	ny kind of
– oral, anal	or vaginal sex.			
Some STDs are fairly comm	non, so there's nothing to be er	nbarrassed about if you ever get one. So	me STDs do not cause any	so
many people who have an	STD don't know it and might _	it. For those who do have s	ymptoms, they may experience sorene	ess or
around the	genitals; bumps sores or rashe	s on the genitals; itching, burning or	while urinating; and gir s m	ay get a
vaginal discharge that is di	fferent from normal.			
If you have any of these sy	mptoms, or if you've had	sex, it's important to get tested	l for STDs. STD tests differ. Some are bl	lood
tests, some test urine, and	still others use a swab of saliva	or a discharge. Even if you use	, it's a good idea to get tested for	r STDs at
least once a year. You can g	get an STD test from your	provider or at a family planning	g or STD clinic.	
Some STDs can be	, while others cannot. B	ut, their symptoms can be managed with	medication. Some STDs, if left untreat	ed, can
		nber, once a person becomes sexually ac		
	get tested for STDs regularly.			
Saltz, Gail, et al. What Are STDs? #F ctChe	ck, Amaze.Org, https://amaze.org/video/what-ard	e-stds-factcheck/. (Accessed October 2, 2019)		
Sexually T	ransm tted	Diseases (STDs		
•		•		
•		ctions or illnesses that can be spread from	•	_
contact. Th	at includes genital-to-	contact, which is contact with per	ises, scrotums, vulvas or vaginas, to an	ıy kind of
– oral, anal	or vaginal sex.			
Some STDs are fairly comm	non, so there's nothing to be er	mbarrassed about if you ever get one. So	me STDs do not cause any	. 50
•		it. For those who do have s		
		s on the genitals; itching, burning or	while urinating; and gir s m	ay get a
vaginal discharge that is di	fferent from normal.			
If you have any of these sy	mptoms, or if you've had	sex, it's important to get tested	l for STDs. STD tests differ. Some are bl	lood
tests, some test urine, and	still others use a swab of saliva	or a discharge. Even if you use	, it's a good idea to get tested for	r STDs at
		provider or at a family planning		
		ut, their symptoms can be managed with		red can
		nber, once a person becomes sexually ac		
		inder, office a person becomes sexually ac	ive, it is important to use	anu
other latex parriers and to	get tested for STDs regularly.			

SEXUALLY TRANSMITTED INFECTIONS AMONG YOUNG AMERICANS

Americans ages 15-24 make up just **27%** of the sexually active population

But account for 50% of the 20M new **STIs** in the U.S. each year



Undiagnosed STIs cause **24,000** women to become infertile each year

HPV

HIV

Syphilis

Total Infections: 820,000 all ages)

2.9 million

14.1 million

776,000

47,500 A ges 13-24

55,400

Gonorrhea

570,000

1 million

Chlamydia

200,000

1.8 million

Diagnosed & reported

Estimated total new infections

Data are cases among youth ages 15-24







GET VACCINATED AGAINST HPV

